



Relevant Kentucky Factsi

Documented Shortage of Primary Care Health Professionals

- According to the Health Resources and Services Administration, Kentucky has 176 total primary care Health Professional Shortage Areas (HPSA).
- Kentucky's population living in a primary care HPSA equals 1,166,737, which represents 26
 percent of the overall population of Kentucky.
- There are 200 licensed audiologists in the Commonwealth of Kentucky who could help.

Hearing Loss Statistics

- Of the more than 36 million Americans who experience hearing loss, 30 percent are 65-74 years old and 47 percent are 75 years old and older.
- Kentucky has 916,000 residents with hearing loss, which represents 20.5 percent of the state population.
- Untreated hearing loss results in a greater occurrence of other physical and mental conditions in elderly patients, substantially reducing the quality of life for older adults and increasing
 Medicare spending. Treated hearing loss enhances the quality of life and reduces health care costs.

National Data on Older Adult Falls

- Every 18 seconds, an older adult is treated in an emergency room for a fall.
- The average cost for a fall injury is \$30,000 and the direct medical cost for falls is \$50 billion annually.
- Among older adults (those 65 or older), falls are the leading cause of injury death. They are also the most common cause of nonfatal injuries and hospital admissions for trauma.
- Elderly falls are preventable as are their associated financial and societal costs. Audiologists reliably diagnose and treat the vestibular disorders and help prevent older adult falls.

Sources

Medicare Beneficiaries by State https://www.kff.org/medicare/state-indicator/total-medicare-beneficiaries/?currentTimeframe=0&sortModel=%7B%22colId%22:%22Location%22,%22sort%22:%22asc%22%7D

 $\label{lem:medicare-spending-by-residence/scurrentTimeframe=0&sortModel=\%7B\%22colld\%22:\%22Location\%22,\%22sort\%22:\%22asc\%22\%7D$

Bureau of Health Workforce, Health Resources and Services Administration (HRSA), U.S. Department of Health & Human Services, <u>Designated Health Professional Shortage Areas Statistics: Designated HPSA Quarterly Summary, as of December 31, 2018</u>.

Audiologists by State https://www.bls.gov/data/

Blackwell, DL and Norris, T. Any Hearing Loss by State: Estimates from the National Health Interview Survey, United States, 2014–2016. National Center for Health Statistics. October 2017. Available from: https://www.cdc.gov/nchs/health-policy/disability.htm.

Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. Accessed August 5, 2016.Burns EB, Stevens JA, Lee RL. The direct costs of fatal and non-fatal falls among older adults—United States. J Safety Res 2016:58.

Safety of audiology direct access for Medicare patients complaining of impaired hearing (Zapala, 2010), http://www.ncbi.nlm.nih.gov/pubmed/20701834;

Screening for vestibular disorders: A study of clinicians' compliance with recommended practices (Polensek, Sterk, Tusa, 2008), http://www.medscimonit.com/download/index/idArt/855748